

# Soils Map




Soils data provided by USDA and NRCS.




State: **Wisconsin**
County: **Green**
Location: **21-3N-8E**
Township: **Mount Pleasant**
Acres: **239.63**
Date: **11/12/2026**

Maps Provided By:
surety® CUSTOMIZED ONLINE MAPPING
© AgriData, Inc. 2025 www.AgriDataInc.com

N W E S

Area Symbol: WI045, Soil Area Version: 27

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Non-Irr Class *c | Irr Class *c | Alfalfa hay Tons | Corn Bu | Corn silage Tons | *n NCCPI Overall | *n NCCPI Corn | *n NCCPI Small Grains | *n NCCPI Soybeans |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 7489A | Marshan silt loam, rarely flooded | 48.46 | 20.3% | | IIw | | | 110 | 18 | 30 | 24 | 15 | 28 |
| 474B | Dickinson sandy loam, 1 to 3 percent slopes | 33.27 | 13.9% | | IIIs | | 4 | 95 | 15 | 52 | 52 | 48 | 36 |
| 251C2 | Northfield loam, 6 to 12 percent slopes, moderately eroded | 24.47 | 10.2% | | IVe | | 2 | 55 | 9 | 44 | 44 | 34 | 28 |
| 108B2 | Tama silt loam, driftless, 2 to 6 percent slopes, moderately eroded | 21.16 | 8.8% | | IIe | | | | | 87 | 87 | 73 | 75 |
| 629A | Ettrick silt loam, 0 to 2 percent slopes, frequently flooded | 17.80 | 7.4% | | VIw | | | | | 52 | 40 | 18 | 52 |
| 478A | Lawler loam, 0 to 2 percent slopes | 16.78 | 7.0% | | IIw | | 3.8 | 110 | 18 | 65 | 65 | 61 | 47 |
| 404A | Dakota loam, 0 to 2 percent slopes | 16.02 | 6.7% | | IIs | | 4.6 | 110 | 19 | 65 | 65 | 58 | 49 |
| 7772A | Marshan loam, rarely flooded | 14.66 | 6.1% | | IIw | | | 110 | 18 | 29 | 25 | 14 | 27 |
| 236C2 | Eleva sandy loam, 6 to 12 percent slopes, moderately eroded | 10.23 | 4.3% | | IIIe | | 3.6 | 75 | 12 | 48 | 48 | 42 | 34 |

Soils data provided by USDA and NRCS.

Maps Provided By:

© AgriData, Inc. 2025 www.AgriDataInc.com

| Code | Soil Description | Acres | Percent of field | Non-Irr Class Legend | Non-Irr Class *c | Irr Class *c | Alfalfa hay Tons | Corn Bu | Corn silage Tons | *n NCCPI Overall | *n NCCPI Corn | *n NCCPI Small Grains | *n NCCPI Soybeans |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 257B2 | Sylvester silt loam, 2 to 6 percent slopes, moderately eroded | 6.79 | 2.8% | | IIe | | 4.7 | 115 | 19 | 66 | 66 | 60 | 57 |
| 246D2 | Elkmound sandy loam, 12 to 20 percent slopes, moderately eroded | 5.54 | 2.3% | | VIe | | 1.8 | 45 | 7 | 34 | 34 | 27 | 20 |
| 404B2 | Dakota loam, 2 to 6 percent slopes, moderately eroded | 5.36 | 2.2% | | IIe | | 4.4 | 105 | 17 | 64 | 64 | 58 | 48 |
| 212B2 | Hixton loam, 2 to 6 percent slopes, moderately eroded | 5.09 | 2.1% | | IIe | | | | | 63 | 63 | 56 | 54 |
| 1246F | Elkmound-Northfield complex, 30 to 60 percent slopes, very rocky | 4.77 | 2.0% | | VIIe | | | | | 10 | 10 | 7 | 3 |
| 606B | Huntsville silt loam, 2 to 6 percent slopes, rarely flooded | 3.41 | 1.4% | | IIe | | 5 | 138 | 23 | 80 | 80 | 73 | 76 |
| 251B2 | Northfield loam, 2 to 6 percent slopes, moderately eroded | 1.99 | 0.8% | | IIIe | | 2.4 | 65 | 10 | 45 | 45 | 35 | 28 |
| 609A | Otter silt loam, frequently flooded | 1.34 | 0.6% | | IIIw | | | 140 | 23 | 38 | 18 | 37 | 18 |
| 154C2 | Lindstrom sandy loam, 6 to 12 percent slopes, moderately eroded | 0.95 | 0.4% | | IIIe | | | | | 78 | 78 | 67 | 73 |
| 305B | Richwood silt loam, 1 to 6 percent slopes | 0.62 | 0.3% | | IIe | | | | | 90 | 89 | 79 | 84 |
| 821B2 | Plainfield loamy sand, 0 to 6 percent slopes, moderately eroded | 0.61 | 0.3% | | IVs | IIIe | 2.5 | 55 | 9 | 31 | 30 | 30 | 24 |
| 301B2 | Pillot silt loam, 2 to 6 percent slopes, moderately eroded | 0.31 | 0.1% | | IIe | | 4.5 | 115 | 19 | 79 | 79 | 70 | 66 |
| | | | **Weighted Average** | | **2.90** | ***-** | **1.9** | **76.3** | **12.5** | ***n 50.1** | ***n 47.7** | ***n 38.9** | ***n 40.1** |

*n: The aggregation method is "Weighted Average using all components"
*c: Using Capabilities Class Dominant Condition Aggregation Method
*- Irr Class weighted average cannot be calculated on the current soils data due to missing data.